

ITP-SB-93-08
February 3, 1993

Wave-type Solutions in the Nonlinear σ -model with the Dilaton

Eugene Tyurin

Institute for Theoretical Physics, SUNY at Stony Brook
E-mail: gene@insti.physics.sunysb.edu

Abstract

We study a class of classical dilaton vacua in string theory that depend on the light-cone variable $z = t \pm x$ and, thus, have wavelike behavior. One of the interesting results is the existence of a solution subclass with perfectly regular space-time geometry, where the string coupling constant can be made arbitrarily small.

Classical solutions in the string theory correspond to conformal field theories — they are represented by the set of background fields that allows the existence of the tree-level world-sheet (sphere in some chosen parameterization). Thus these classical solutions can be called stringy vacua, as loop corrections are identified with the quantum fluctuations of the world-sheet.

In this paper, starting from one particular form of the space-time geometry, we will find the wide class of the wave-type classical solutions that can be made to have singular as well as regular behavior.

We establish our notation. The σ -model action with the dilaton term is

$$S_\sigma = \frac{1}{4\pi\alpha'} \int d^2\sigma \sqrt{h} \left[h^{ab} \partial_a X^m \partial_b X^n g_{mn}(X^r) - \frac{\alpha'}{2} \phi(X^r) R^{(2)} \right], \quad (1)$$

where h^{ab} is the world-sheet metric, g_{mn} is the metric on the D-dimensional space-time with the coordinates X^r , $\phi(X^r)$ denotes the dilaton field, and $R^{(2)}$ is the world-sheet curvature scalar.

To lowest order in α' , the β -functions that should be equal to zero to insure both conformal invariance and ultraviolet finiteness of the action (1), are

$$\beta_{mn}^G = R_{mn} - \nabla_m \nabla_n \phi = 0 \quad (2)$$

$$\beta^\Phi = R + (\nabla\phi)^2 - 2\nabla^2\phi + \Lambda = 0, \quad (3)$$

where $\Lambda = (D - D_{crit})/3\alpha'$ plays the role of “cosmological constant” in the low-energy effective action

$$S_{eff} = \int d^D X \sqrt{g} e^{-\phi} [R^{(D)} - (\nabla\phi)^2 + \Lambda] \quad (4)$$

Note that in general, the dilaton allows us to formulate the theory with the target space dimensionality $D \neq D_{crit}$, because without it, the field equations (2) and (3) turn into

$$\begin{cases} R_{mn} = 0 \\ R + \Lambda = 0 \end{cases} \implies \Lambda = 0 \Leftrightarrow D = D_{crit}$$

In this paper we do not discuss other possible background fields: axion and tachyon, because (1) the axion can be easily generated by duality transformations [1]; (2) the tachyon is absent in superstrings.

Recently a lot of attention has been given to the study of conformal field theory with the geometry of the target space

$$ds^2 = \mp dt^2 + a^2(t) d\theta^2 + \sum_{i=1}^{D-2} d\xi_i^2, \quad (5)$$

and the dilaton $\phi(t)$, where t is a Minkowskian or Euclidean time. In this case β -functions (2), (3) have the following form:

$$-\beta_{tt}^G = \frac{\ddot{a}}{a} + \ddot{\phi} = 0 \quad (6)$$

$$\beta_{\theta\theta}^G = \frac{\ddot{a}}{a} + \frac{\dot{a}}{a}\dot{\phi} = 0 \quad (7)$$

$$\beta^\Phi = 2\frac{\ddot{a}}{a} - (\dot{\phi})^2 + 2\ddot{\phi} + 2\frac{\dot{a}}{a}\dot{\phi} + \Lambda = 0, \quad (8)$$

where as usual “dot” $\equiv \partial/\partial t$. It turns out that there are only four nontrivial solutions for the equations (6)–(8):

$$\begin{aligned} a(t) &= \tan(\omega t), \cot(\omega t), \tanh(\omega t), \coth(\omega t) \\ \phi(t) &= \phi_0 + \ln \bar{a}^2(t), \text{ where} \\ \bar{a}(t) &= \sin(\omega t), \cos(\omega t), \sinh(\omega t), \cosh(\omega t) \text{ correspondingly} \end{aligned} \quad (9)$$

As one can notice these solutions describe the black-hole type singularity (cf. [2, 3]) and the choice of D determines the value of ω ($a(t) = 1/t$, $\phi(t) = \phi_0 + \ln t^2$ is possible only in the case of $\Lambda = 0$ [3]).

An obvious generalization of the metric (5) (see, for example, paper [4] and references therein) is

$$ds^2 = -dt^2 + dx^2 + a^2(t)d\theta_1^2 + b^2(x)d\theta_2^2 + \sum_{i=1}^{D-4} d\xi_i^2 \quad (10)$$

But it is still essentially two-dimensional. Another variant: curved inclusion of the compactified coordinates discussed in the work [3]

$$\begin{aligned} ds^2 &= -dt^2 + \sum_{i=1}^{D-1} [2\pi r_i(t)]^2 dX_i^2, \\ X^i &\equiv X^i + 1 \end{aligned} \quad (11)$$

doesn't bring us any drastically distinct solutions for the “radii” $r_i(t)$.

For these reasons we consider the following geometry of the space-time:

$$ds^2 = -dt^2 + dx^2 + a^2(t, x)d\theta_1^2 + b^2(t, x)d\theta_2^2 + \sum_{i=1}^{D-4} d\xi_i^2 \quad (12)$$

$$\phi = \phi(t, x)$$

Here we keep the main features of the equations (5), (10) and (11) while looking for the nontrivial both in t and x functional dependence of the metric coefficients and the dilaton field.

Equations (2) have now the following form:

$$-\beta_{tt}^G = \frac{\ddot{a}}{a} + \frac{\ddot{b}}{b} + \ddot{\phi} = 0, \quad (13)$$

$$-\beta_{xx}^G = \frac{a''}{a} + \frac{b''}{b} + \phi'' = 0, \quad (14)$$

$$-\beta_{tx}^G = \frac{\dot{a}'}{a} + \frac{\dot{b}'}{b} + \dot{\phi}' = 0, \quad (15)$$

$$\beta_{\theta_1\theta_1}^G = \frac{\ddot{a}}{a} - \frac{a''}{a} + \frac{a'b'}{ab} - \frac{\dot{a}\dot{b}}{ab} + \frac{\dot{a}}{a}\dot{\phi} - \frac{a'}{a}\phi' = 0, \quad (16)$$

$$\beta_{\theta_2\theta_2}^G = \frac{\ddot{b}}{b} - \frac{b''}{b} + \frac{a'b'}{ab} - \frac{\dot{a}\dot{b}}{ab} + \frac{\dot{b}}{b}\dot{\phi} - \frac{b'}{b}\phi' = 0, \quad (17)$$

where “prime” $\equiv \partial/\partial x$. One can notice that the above system includes the two types of partial differential equations and allows the following compact representation:

$$\left(\frac{\partial^2 \phi}{\partial X^i \partial X^j} + \overrightarrow{\partial}_i \overrightarrow{\partial}_j \right) C^\kappa(t, x) C^\lambda(t, x) = 0 \quad \kappa \neq \lambda, \quad (18)$$

$$C^\kappa(t, x) \eta^{ij} \left[\overrightarrow{\partial}_i - \overleftarrow{\partial}_i + \frac{\partial \phi}{\partial X^i} \right] \overrightarrow{\partial}_j C^\lambda(t, x) = 0 \quad \kappa \neq \lambda, \quad (19)$$

where $C^{\kappa,\lambda}(t, x)$ denotes $a(t, x)$ or $b(t, x)$, $X^i = \{t, x\}$ and $\eta^{ij} = \text{diag}\{-1, 1\}$. Unfortunately we do not know the general solution for this problem and, though the proposed geometry is an obvious generalization of (5), (10), (11), such “obvious” assumptions as $a, b = f(t)g(x)$ or $[f(t) + g(x)]^p$ do not work unless they accidentally fall in the solution class that will be introduced below.

Suppose a, b, ϕ to depend only on the single light-cone variable $z = t + x$ or $\tilde{z} = t - x$. All the subsequent formulae do not depend on the choice of z or $\tilde{z}$, so for the sake of clarity we will speak in the terms of z . With the above ansatz equations (19) are satisfied identically for any a, b, ϕ and the three equations denoted as (18) are equivalent to each other, and are just

$$\frac{a''(z)}{a(z)} + \frac{b''(z)}{b(z)} + \phi''(z) = 0,$$

where “prime” from now on means $\partial/\partial z$. The only component of the Ricci tensor that is not equal to zero identically is

$$R_{zz} = -\frac{a''}{a} - \frac{b''}{b}, \quad (20)$$

hence $R = R_a{}^a \equiv 0$. One can also check that both $\nabla^2 \phi \equiv 0$ and $(\nabla \phi)^2 \equiv 0$. Because of these cancellations, equation (3) is equivalent to the condition $\Lambda = 0 \Leftrightarrow D = D_{crit}$ and this is distinct from the previously discussed cases.

So the full system of the lowest order β -function equations we should satisfy is

$$\frac{a''}{a} + \frac{b''}{b} + \phi'' = 0, \quad (21)$$

$$\Lambda = 0 \quad (22)$$

Since the equations (18) and (6) are similar to each other, one can employ as the wave-type solutions all of the options (9) which should be identified now with moving singularities.

Generally, we are not interested in the higher orders in α' , because in the supersymmetric theory one can always make the quasi-classical approximation to be exact, but it is nice to notice that the second-order corrections to the β -functions that are proportional to $\delta\beta_{mn} = \alpha' R_{mabc} R_n{}^{abc}$ are identically equal to zero in our case.

Another interesting feature of the solution class we have found is that we can always change the constant and linear powers of the dilaton field (there are no first powers of the ordinary derivatives involved).

Of course, since we have only one equation for three functions, there is a tremendous freedom of choice. For example, after fixing the dilaton field, we obtain the following equations for $a(z), b(z)$ with an absolutely arbitrary function $f(z)$:

$$\begin{aligned} \frac{a''}{a} + \phi'' - f &= 0, \\ \frac{b''}{b} + f &= 0, \end{aligned} \quad (23)$$

that have the general form of $y''(x) + w(x)y(x) = 0$. This type of differential equations has been thoroughly studied in mathematics (for a review see, for example, [5]) and here we will just say that the existence of a finite or infinite number of zeros for $y(x)$ (i.e. metric irregularities in our terminology) is guaranteed for a wide class of $w(x)$.

Since the ultimate purpose of the model-building approach that we employ here is to obtain something that seems to be a reasonable choice of the classical vacuum in the string theory, it seems to be interesting to find solutions with a “perfect behavior”. Recall that, since we work in

the perturbation theory with the effective string coupling constant $g = \exp(\phi/2)$, and the dilaton $\phi(z)$ can always be shifted by some finite constant ϕ_0 , g can always be made arbitrarily (but, in general, not infinitely) small if $\phi(z) \neq +\infty$ for all z . Additionally we will require the target space metric g_{mn} to be regular everywhere except perhaps light-cone infinity. The following statement can point to some of these “nice” solutions.

Theorem Let $a(z) \in C^2(\mathbb{R})$ be a symmetric, monotonically non-increasing for $z > 0$ function and let

$$\lim_{z \rightarrow 0} a(z) = c_1 < \infty \quad \lim_{z \rightarrow +\infty} a(z) = c_2 \geq 0, \text{ then the dilaton}$$

$$\phi_a(z) = - \int \int \frac{a''}{a} dz dz \rightarrow -\infty \text{ as } z \rightarrow +\infty$$

and $\phi_a(z) < +\infty$ for all z .

Proof First of all, note that from the condition $a(z) = a(-z)$ and from the continuity of $a'(z)$ follows that there exists a point $\zeta > 0$ such that

$$a''(z) \begin{cases} < 0 & z < \zeta \\ > 0 & z > \zeta \end{cases}, \quad \text{then } \phi_a'' = -\frac{a''}{a} \begin{cases} > 0 & z < \zeta \\ < 0 & z > \zeta \end{cases}$$

It is clear that ϕ_a'' cannot be positive for $z > \zeta$, because $\lim_{z \rightarrow +\infty} a''(z) = -0$. On the next step, there is such $\gamma > \zeta$, that

$$\phi_a' = \int dz \phi_a'' \begin{cases} > 0 & z < \gamma \\ < 0 & z > \gamma \end{cases}$$

Again, ϕ_a' cannot be positive for $z > \gamma$, because of the infiniteness of the interval where $\phi_a'' < 0$. Finally, we find that $\lim_{z \rightarrow +\infty} \phi_a(z) = -\infty$, because $\phi_a' < 0$ for $z > \gamma$. **Q.E.D.**

Note that if we consider monotonically increasing $a(z)$ for $z > 0$ then $\lim_{z \rightarrow +\infty} \phi_a(z) = +\infty$. Unfortunately it is impossible to analyse oscillating, non-monotonic functions in the same fashion. But, with the theorem proven above, one can pick a lot of metrics that contain this “nice” dilaton background (cf. (23): of course, we assume here that $f(z) = -b''/b = \phi_b''$).

We have discussed the class of the classical dilaton solutions to the critical string theory that allows a wave-packet interpretation. These solutions allow both singularities and finite, smooth backgrounds. Since for the space-time geometry we have considered, there are some miraculous cancellations in the β -function equations, it may be interesting to investigate this topic for some more general initial assumptions. The vacua presented in this paper can also be found in [6], where the metrics of the form (11), but in the light-cone variables were studied.

I would like to express my deep gratitude to M. Roček for suggesting this problem to me and for his advice.

References

- [1] A. Giveon and M. Roček *Generalized Duality in Curved String-Backgrounds* preprint IASSNS-HEP-91/84 and ITP-SB-91-67
- [2] E. Witten *String Theory and Black Holes* Phys. Rev. **D44** (1991), 314
- [3] M. Mueller *Rolling Radii and a Time-Dependent Dilaton* Nucl. Ph. **B337** (1990), 37
- [4] A. Giveon and A. Pasquinucci *On Cosmological String Backgrounds with Toroidal Isometries* preprint IASSNS-HEP-92/55 and PUPT-1337
- [5] E. Kamke *Differentialgleichungen. Lösungsmethoden und Lösungen.* 1959 Leipzig, Akademische Verlagsgesellschaft
- [6] R. Rudd *Compactification Propagation* Nucl. Ph. **B352** (1991), 489
- [7] E. Bergshoeff, R. Kallosh and T. Ortin *Supersymmetric String Waves* preprint SU-ITP-92-30 and UG-10/92